arXiv:1001.1220v2 [math.AC] 23 Mar 2011

JUMPING NUMBERS AND ORDERED TREE STRUCTURES ON THE DUAL GRAPH

EERO HYRY AND TARMO JÄRVILEHTO

Dedicated to the memory of Professor Olli Jussila

Abstract. Let R be a two-dimensional regular local ring having an algebraically closed residue field and let $\mathfrak{a}$ be a complete ideal of finite colength in R. In this article we investigate the jumping numbers of $\mathfrak{a}$ by means of the dual graph of the minimal log resolution of the pair $(X, \mathfrak{a})$. Our main result is a combinatorial criterium for a positive rational number ξ to be a jumping number. In particular, we associate to each jumping number certain ordered tree structures on the dual graph.

1. Introduction

Multiplier ideals have in recent years emerged as an important tool in algebraic geometry. Given a closed subscheme of a smooth complex variety, there is a nested sequence of multiplier ideals parametrized by the positive rational numbers. A jumping number is a value of the rational parameter at which the multiplier ideal makes a jump. Jumping numbers form a discrete set of invariants, which contains important information about the singularities of the subscheme in question.

Jumping numbers are defined by using an embedded resolution of the subcheme. They depend on the exceptional divisors appearing in the resolution. The purpose of this article is to look at jumping numbers from the point of view of the combinatorics of exceptional divisors in the case of a smooth surface. In particular, we associate to each jumping number certain ordered tree structures on the dual graph of the resolution. These structures generalize the one discovered by Veys in [11] while he was studying poles of the topological zeta function

To describe our work in more detail, let $\mathfrak{a}$ be a complete ideal of finite colength in a two-dimensional regular local ring R having an algebraically closed residue field. Let $X \longrightarrow \operatorname{Spec} R$ be a minimal log resolution of the pair $(R, \mathfrak{a})$. Let D be the divisor on X such that $\mathcal{O}_X(-D) = \mathfrak{a}\mathcal{O}_X$. Let $E_1, \ldots, E_N$ be the exceptional divisors. Recall that a divisor $F = f_1E_1 + \ldots + f_NE_N$ is called antinef if $F \cdot E_\gamma \leq 0$ for all $\gamma = 1, \ldots, N$, where $F \cdot E_\gamma$ denotes the intersection product. Our starting point is the observation made in [7] that jumping numbers of $\mathfrak{a}$ can be parametrized by the antinef divisors. More precisely, the jumping number corresponding to F is

$$\xi_F := \min_{\gamma} \frac{f_\gamma + k_\gamma + 1}{d_\gamma},$$

where $D = d_1E_1 + \ldots + d_NE_N$ and $K = k_1E_1 + \ldots + k_NE_N$ is the canonical divisor.

Let Γ denote the dual graph of X. Recall that the vertices of the dual graph correspond to the exceptional divisors and that two vertices are adjacent if and only if the corresponding exceptional divisors intersect. The above considerations motivate us to investigate the function $\lambda(f_\gamma, \gamma)$ on $|\Gamma|$, where

$$\lambda(a, \gamma) := \frac{a + k_\gamma + 1}{d_\gamma}$$

for any $a \in \mathbb{Z}$ and $\gamma \in |\Gamma|$. By the above the minimum value of $\lambda(f_\gamma, \gamma)$ is now the jumping number $\xi = \xi_F$. We call the set

$$S_F := \{\gamma \in |\Gamma| \mid \lambda(f_\gamma, \gamma) = \xi\}$$

the support of ξ with respect to the antinef divisor F.

Our main result Theorem 4.3 is a combinatorial criterium for a positive rational number ξ to be a jumping number. We observe that it is possible to choose the divisor F in such a way that $\lambda(f_\gamma, \gamma)$ strictly increases along every path away from the support. By assigning the number $\lambda(f_\gamma, \gamma)$ to each vertex γ, we can make the dual graph an ordered tree. An end of S_F must either have at least three adjacent vertices or correspond to some Rees valuation of the ideal. Moreover, one may assume that S_F is a chain such that the vertices corresponding to Rees valuations do not occur at the non-ends of S_F.

We also want to understand how the so called contributing divisors arise. The notion of contribution to a jumping number by a divisor was introduced by Smith and Thompson in [9], and the investigation has then been continued by Tucker in [10]. It turns out in Theorem 4.16 that every critically contributing divisor, in the sense of Tucker, is of the type $\sum_{\gamma \in S_F} E_\gamma$. However, the converse is not true. Therefore we give in Theorem 4.19 a necessary and sufficient condition for a reduced divisor to be a critically contributing one.

Using Theorem 4.3 we will also show in Corollary 4.10 that given a vertex with at least three adjacent vertices or a vertex corresponding to a Rees valuation of the ideal, there is always a jumping number supported exactly at this vertex. Note that a support of a jumping number always contains vertices of this type. Moreover, by means of Theorem 4.3 we can in Proposition 4.12 construct from a given jumping number certain new jumping numbers having the same support as the original one.

Our main technical tool is Lemma 3.6 which helps us to construct suitable antinef divisors F. This is inspired by the work of Loeser and

Veys concerning the numerical data associated to the exceptional divisors of the resolution (see [6] and [11]).

Finally, we would like to refer to the book of Favre and Jonsson ([3]) for related topics. It would be interesting to know whether our results can be interpreted in their 'tree language'.

2. Preliminaries

We begin by fixing notation and recalling some basic facts from the Zariski-Lipman theory of complete ideals. For more details, we refer to [4], [1], [5] and [7].

Throughout this article R denotes a regular local ring of dimension two having an algebraically closed residue field. Let $\mathfrak{a}$ be a complete ideal of finite colength in R. Let $\pi\colon X \to \operatorname{Spec}(R)$ be a minimal principalization of $\mathfrak{a}$. Then X is a regular scheme and $\mathfrak{a}\mathcal{O}_X = \mathcal{O}_X(-D)$ for an effective Cartier divisor D. Note that π is a log resolution of the pair $(\operatorname{Spec} R, \mathfrak{a})$, i.e., the divisor $D + \operatorname{Exc}(\pi)$ has simple normal crossing support, where $\operatorname{Exc}(\pi)$ denotes the sum of the exceptional divisors of π.

The morphism π is a composition of point blowups of regular schemes

$$\pi : X = X_{N+1} \xrightarrow{\pi_N} \cdots \xrightarrow{\pi_2} X_2 \xrightarrow{\pi_1} X_1 = \operatorname{Spec} R,$$

where $\pi_\mu : X_{\mu+1} \to X_\mu$ is the blowup of X_μ at a closed point $x_\mu \in X_\mu$ for every $\mu = 1, \ldots, N$. Let E_μ and E_μ^*, respectively, be the strict and total transform of the exceptional divisor $\pi_\mu^{-1}\{x_\mu\}$ on X for every $\mu = 1, \ldots, N$. We denote by v_μ the discrete valuation associated to the discrete valuation ring $\mathcal{O}_{X,E_\mu}$, in other words, v_μ is the $\mathfrak{m}_{X_\mu,x_\mu}$-adic order valuation.

Recall that a point x_μ is said to be infinitely near to a point x_ν, if the projection $X_\mu \to X_\nu$ maps x_μ to x_ν. This relation gives a partial order on the set $\{x_1, \ldots, x_N\}$. A point x_μ is *proximate* to the point x_ν, denoted by $\mu \succ \nu$, if and only if x_μ lies on the strict transform of $\pi_\nu^{-1}\{x_\nu\}$ on X_μ. Following [1, Definition-Lemma 1.5], the *proximity matrix* is

$$P := (p_{\mu,\nu})_{N\times N}, \text{ where } p_{\mu,\nu} = \begin{cases} 1, & \text{if } \mu = \nu; \\ -1, & \text{if } \mu \succ \nu; \\ 0, & \text{otherwise.} \end{cases}$$

Note that this is the transpose of the proximity matrix given in [4, p. 6]. We set $Q = (q_{\mu,\nu})_{N\times N} := P^{-1}$. The equation $PQ = 1$ immediately gives the formula

$$q_{\mu,\nu} = \sum_{\mu \succ \rho} q_{\rho,\nu} + \delta_{\mu,\nu}. \tag{1}$$

If x_μ is infinitely near to x_ν, then $q_{\mu,\nu} > 0$ while $q_{\mu,\nu} = 0$ otherwise. Clearly $q_{\mu,\mu} = 1$ for all $\mu = 1, \ldots, N$.

We denote by Γ the *dual graph* associated to our principalization. It is well known that Γ is a tree. Let $|\Gamma|$ be the corresponding set of vertices.

Recall that there is a vertex ν corresponding to each exceptional divisor E_ν weighted by the number $w_\Gamma(\nu) := -E_\nu^2$. Note that

$$w_\Gamma(\nu) = 1 + \#\{\mu \in |\Gamma| \mid \mu \succ \nu\}.$$

Two vertices μ and ν are called *adjacent* if they can be joined by an edge. This is the case if and only if the corresponding exceptional divisors E_μ and E_ν intersect. We write $\mu \sim \nu$. Then either $\mu \prec \nu$ or $\mu \succ \nu$. Suppose, for example, that $\mu \succ \nu$. Then $\nu \sim \mu$ in fact means that x_μ is a maximal element in the set of infinitely near points proximate to x_ν. The *valence* $v_\Gamma(\nu)$ of a vertex ν means the number of vertices adjacent to it. If $v_\Gamma(\nu) \geq 2$, then ν is called a *star.* A vertex τ with $v_\Gamma(\tau) = 1$ is an *end.* The *distance* between two vertices $\mu, \nu \in |\Gamma|$ is defined as

$$d(\nu, \mu) := \min\{r \mid \nu = \nu_0 \sim \cdots \sim \nu_r = \mu, \text{ where } \nu_0, \ldots, \nu_r \in |\Gamma|\}.$$

Furthermore, if $S \subset |\Gamma|$, we set

$$d(\nu, S) := \min\{d(\nu, \mu) \mid \mu \in S\}.$$

If $d(\nu, S) = 1$, then we write $\nu \sim S$.

We consider the lattice $\Lambda := \mathbb{Z}E_1 \oplus \ldots \oplus \mathbb{Z}E_N$ of exceptional divisors on X. The lattice Λ has two other convenient bases besides $\{E_\mu \mid \mu = 1, \ldots, N\}$, namely $\{E_\mu^* \mid \mu = 1, \ldots, N\}$ and $\{\widehat{E}_\mu \mid \mu = 1, \ldots, N\}$, where $E_\mu \cdot \widehat{E}_\nu = -\delta_{\mu,\nu}$ for $\mu, \nu = 1, \ldots, N$. For any $G \in \Lambda$, we write

$$G = g_1E_1 + \ldots + g_NE_N = g_1^*E_1^* + \ldots + g_N^*E_N^* = \widehat{g}_1\widehat{E}_1 + \ldots + \widehat{g}_N\widehat{E}_N.$$

The following base change formulas now hold:

$$g^* = gP^{\mathrm{T}} \text{ and } \widehat{g} = gP^{\mathrm{T}}P = g^*P, \tag{2}$$

where g, g^* and $\widehat{g}$ denote row vectors in $\mathbb{Z}^n$. Here $(P^{\mathrm{T}}P)_{\mu,\nu} = -E_\mu \cdot E_\nu$. In particular, note the formulas

$$\widehat{g}_\mu = g_\mu^* - \sum_{\nu \succ \mu} g_\nu^* = w_\Gamma(\gamma)g_\mu - \sum_{\nu \sim \mu} g_\nu \quad (\mu = 1, \ldots, N). \tag{3}$$

The *support* of a divisor $G \in \Lambda$ is $|G| := \{\gamma \in |\Gamma| \mid g_\gamma \neq 0\}$.

Recall that a divisor $F \in \Lambda$ is *antinef* if $\widehat{f}_\nu = -F \cdot E_\nu \geq 0$ for all $\nu = 1, \ldots, N$. Equivalently, the *proximity inequalities*

$$f_\mu^* \geq \sum_{\nu \succ \mu} f_\nu^* \quad (\mu = 1, \ldots, N) \tag{4}$$

hold. Note that they can also be expressed in the form

$$w_\Gamma(\mu)f_\mu \geq \sum_{\nu \sim \mu} f_\nu \quad (\mu = 1, \ldots, N). \tag{5}$$

In fact, if $F \neq 0$ is antinef, then also $f_\nu > 0$ for all $\nu = 1, \ldots, N$. There is a one to one correspondence between the antinef divisors in Λ and the complete ideals of finite colength in R generating invertible $\mathcal{O}_X$-sheaves, given by $F \leftrightarrow \Gamma(X, \mathcal{O}_X(-F))$. For a divisor $G \in \Lambda$, there exists a minimal one among the antinef divisors F satisfying $F \geq G$. This is called the *antinef closure* of G and denoted by $G^{\sim}$. We have

$$\Gamma(X, \mathcal{O}_X(-G)) = \Gamma(X, \mathcal{O}_X(-G^{\sim}))$$

for any divisor $G \in \Lambda$.

Recall that an ideal is called *simple* if it cannot be expressed as a product of two proper ideals. By the famous result of Zariski, every complete ideal factorizes uniquely into a product of simple complete ideals. More precisely,

$$\mathfrak{a} = \mathfrak{p}_1^{\widehat{d}_1} \cdots \mathfrak{p}_N^{\widehat{d}_N},$$

where $\mathfrak{p}_\mu \subset R$ denotes the simple complete ideal of finite colength corresponding to the exceptional divisor E_μ and $\widehat{d}_\mu > 0$ if and only if v_μ is a Rees valuation of $\mathfrak{a}$. We have $\mathfrak{p}_\mu \mathcal{O}_X = \mathcal{O}_X(-\widehat{E}_\mu)$ so that $\mathfrak{p}_\mu = \Gamma(X, \mathcal{O}_X(-\widehat{E}_\mu))$. By (2)

$$\widehat{E}_\mu = \sum_\nu q_{\mu,\nu} E_\nu^* = \sum_{\nu,\rho} q_{\nu,\rho} q_{\mu,\rho} E_\nu. \tag{6}$$

In particular, we observe the *reciprocity formula*

$$v_\nu(\mathfrak{p}_\mu) = \sum_\rho q_{\nu,\rho} q_{\mu,\rho} = v_\mu(\mathfrak{p}_\nu) \quad (\mu, \nu = 1, \ldots, N). \tag{7}$$

For $\mu \neq \nu$, the proximity inequalities now become equalities

$$q_{\mu,\nu} = \sum_{\rho \succ \nu} q_{\mu,\rho}.$$

We will next recall the definition of jumping numbers. A general reference for jumping numbers is the fundamental article [2]. Recall first that the *canonical divisor* is $K = \sum_\nu E_\nu^*$. The formulas (2) now give

$$k_\nu = \sum_\mu q_{\nu,\mu} \quad \text{and} \quad \widehat{k}_\nu = E_\nu^2 + 2 \quad (\nu = 1, \ldots, N). \tag{8}$$

By (3) we then obtain an important relation

$$w_\Gamma(\nu) k_\nu = 2 - w_\Gamma(\nu) + \sum_{\mu \sim \gamma} k_\mu \quad (\nu = 1, \ldots, N). \tag{9}$$

For a nonnegative rational number ξ, the *multiplier ideal* $\mathcal{J}(\mathfrak{a}^\xi)$ is defined to be the ideal

$$\mathcal{J}(\mathfrak{a}^\xi) := \Gamma\left(X, \mathcal{O}_X\left(K - \lfloor \xi D \rfloor\right)\right) \subset R,$$

where $\lfloor \xi D \rfloor$ denotes the integer part of ξD. It is now known that there is an increasing discrete sequence

$$0 = \xi_0 < \xi_1 < \xi_2 < \cdots$$

of rational numbers ξ_i characterized by the properties that $\mathcal{J}(\mathfrak{a}^\xi) = \mathcal{J}(\mathfrak{a}^{\xi_i})$ for $\xi \in [\xi_i, \xi_{i+1})$, while $\mathcal{J}(\mathfrak{a}^{\xi_{i+1}}) \subsetneq \mathcal{J}(\mathfrak{a}^{\xi_i})$ for every i. The numbers $\xi_1, \xi_2, \ldots,$ are called the *jumping numbers* of $\mathfrak{a}$. Note that contrary to [2, Definition 1.4], we don't consider 0 as a jumping number. Clearly, this is no restriction. The following Proposition 2.1, which is fundamental for the rest of this article, results from [7, Proposition 6.7 and Proposition 7.2].

Proposition 2.1. *Let R be a two-dimensional regular local ring and let $\mathfrak{a} \subset R$ be a complete ideal of finite colength. Then ξ is a jumping number of $\mathfrak{a}$ if and only if there exists an antinef divisor $F \in \Lambda$ such that*

$$\xi = \xi_F := \min_{\nu} \frac{f_\nu + k_\nu + 1}{d_\nu}.$$

Moreover, if $\mathfrak{b}$ is the complete ideal corresponding to F, then

$$\xi = \inf\{c \in \mathbb{Q}_{>0} \mid \mathcal{J}(\mathfrak{a}^c) \not\supseteq \mathfrak{b}\}.$$

Notation 2.2. *We write for any integer a and for any vertex ν*

$$\lambda(a, \nu) := \frac{a + k_\nu + 1}{d_\nu}$$

and call the set

$$S_F := \{\nu \in |\Gamma| \mid \lambda(f_\nu, \nu) = \xi\}$$

the support of the jumping number ξ with respect to the antinef divisor F.

3. RELATIONS BETWEEN NUMERICAL DATA ASSOCIATED TO EXCEPTIONAL DIVISORS

Let ξ be a jumping number of the ideal $\mathfrak{a}$. In order to define an ordered tree structure on the dual graph as described in the introduction, we must be able to construct an antinef divisor F such that $\xi = \xi_F$ and that $\lambda(f_\gamma, \gamma)$ increases along every path away from the support S_F. Proceeding inductively, suppose that we are given a vertex γ and a vertex $\eta_1 \sim \gamma$ such that $d(\gamma, S_F) > d(\eta_1, S_F)$. Suppose, furthermore, that numbers f_γ and f_{η_1} have been defined in such a way that $\lambda(f_\gamma, \gamma) > \lambda(f_{\eta_1}, \eta_1)$. The key issue is to find for vertices $\eta_1 \neq \eta \sim \gamma$ suitable numbers f_η with the property that $\lambda(f_\eta, \eta) > \lambda(f_\gamma, \gamma)$. This problem will be addressed in Lemma 3.5, which is the main result of this section. Other details of the above construction will be postponed till Lemma 4.1 in the next section.

We begin with the following lemma:

Lemma 3.1. *Let $F \in \Lambda$ be a divisor. Let $\gamma \in |\Gamma|$ be a vertex such that $\widehat{f_\gamma} \geq 0$ and $\xi := \lambda(f_\gamma, \gamma) \leq \lambda(f_\eta, \eta)$ when $\eta \sim \gamma$. Then*

$$\xi \widehat{d_\gamma} \geq \widehat{f_\gamma} - \upsilon_\Gamma(\gamma) + 2.$$

In particular, this implies the following:

a) *If $\upsilon_\Gamma(\gamma) \leq 2$ and $\widehat{d_\gamma} = 0$, then there are exactly two vertices η adjacent to γ and $\lambda(f_\eta, \eta) = \xi$ for both of those. Furthermore $\widehat{f_\gamma} = 0$.*

b) *If* $v_\Gamma(\gamma) = 1$*, then* $\xi \widehat{d}_\gamma \geq \widehat{f}_\gamma + 1$*. Especially,* $\widehat{d}_\gamma > 0$*.*

Proof. By the formulas (3) and (9) we obtain

$$\xi = \frac{w_\Gamma(\gamma)(f_\gamma + k_\gamma + 1)}{w_\Gamma(\gamma) d_\gamma} = \frac{\sum_{\eta\sim\gamma}(f_\eta + k_\eta + 1) + \widehat{f}_\gamma - v_\Gamma(\gamma) + 2}{\sum_{\eta\sim\gamma} d_\eta + \widehat{d}_\gamma}.$$

Since

$$\xi \leq \frac{f_\eta + k_\eta + 1}{d_\eta},$$

we must have $\xi \widehat{d}_\gamma \geq \widehat{f}_\gamma - v_\Gamma(\gamma) + 2$.

When $v_\Gamma(\gamma) \leq 1$, this immediately gives $\widehat{d}_\gamma > 0$. Then suppose that $\widehat{d}_\gamma = 0$ and $v_\Gamma(\gamma) = 2$. Now $\widehat{f}_\gamma - v_\Gamma(\gamma) + 2 \leq 0$ is possible only if

$$\widehat{f}_\gamma = v_\Gamma(\gamma) - 2 = 0.$$

But then

$$\xi = \frac{\sum_{\eta\sim\gamma}(f_\eta + k_\eta + 1)}{\sum_{\eta\sim\gamma} d_\eta},$$

which implies that $\xi = \lambda(f_\eta, \eta)$ for both $\eta \sim \gamma$. □

For any two vertices $\nu, \gamma \in |\Gamma|$, set

$$\alpha_{\gamma,\nu} := k_\nu + 1 - \frac{k_\gamma + 1}{d_\gamma} d_\nu = d_\nu(\lambda(0,\nu) - \lambda(0,\gamma)).$$

The numbers $\alpha_{\gamma,\nu}$ were first investigated by Loeser in [6] in the case of an embedded resolution of a curve. Van Proeyen and Veys generalized his results to the ideal case in [8]. The following Lemma 3.2 and Lemma 3.3 are due to them ([8, Proposition 3.1 and Corollary 3.2]). Since we have slightly modified the statements, we include the proofs here for the convenience of the reader. Moreover, working in a more algebraic context, we also prefer to prove these results directly without utilizing the results of Loeser.

Lemma 3.2. *If* $\gamma \in |\Gamma|$*, then*

$$\sum_{\nu\sim\gamma} \alpha_{\gamma,\nu} = v_\Gamma(\gamma) - 2 + \widehat{d}_\gamma \frac{k_\gamma + 1}{d_\gamma}.$$

Proof. By the relation (9)

$$\sum_{\nu\sim\gamma} k_\nu = w_\Gamma(\gamma) k_\gamma + w_\Gamma(\gamma) - 2$$

and by the formula (3)

$$\sum_{\nu\sim\gamma} d_\nu = w_\Gamma(\gamma) d_\gamma - \widehat{d}_\gamma.$$

It follows that

$$\begin{aligned}\sum_{\nu\sim\gamma}\alpha_{\gamma,\nu} &= \sum_{\nu\sim\gamma}k_\nu + v_\Gamma(\gamma) - \sum_{\nu\sim\gamma}d_\nu\frac{k_\gamma+1}{d_\gamma}\\ &= w_\Gamma(\gamma)\,(k_\gamma+1) + v_\Gamma(\gamma) - 2 - \left(w_\Gamma(\gamma)d_\gamma - \widehat{d}_\gamma\right)\frac{k_\gamma+1}{d_\gamma}\\ &= v_\Gamma(\gamma) - 2 + \widehat{d}_\gamma\frac{k_\gamma+1}{d_\gamma}.\end{aligned}$$

□

Lemma 3.3. *Let $\gamma \in |\Gamma|$.*

a) *For every $\eta\sim\gamma$, either*

$$|\alpha_{\gamma,\eta}| < 1,$$

or $\widehat{d}_\gamma = 0$ and η is the only vertex adjacent to γ, in which case $\alpha_{\gamma,\eta} = -1$.

b) *For all except at most one $\eta\sim\gamma$, $\alpha_{\gamma,\eta}\geq 0$. More precisely, if $\alpha_{\gamma,\eta}\leq 0$ for some $\eta\sim\gamma$, then $\alpha_{\gamma,\nu}>0$ for all $\eta\neq\nu\sim\gamma$, unless $\alpha_{\gamma,\eta}=0$, $\widehat{d}_\gamma=0$ and there are only two vertices $\eta'\sim\gamma\sim\eta$, in which case also $\alpha_{\gamma,\eta'}=0$.*

Proof. a) Consider the sequence of point blowups

$$\pi : X = X_{N+1} \xrightarrow{\pi_N} \cdots \xrightarrow{\pi_2} X_2 \xrightarrow{\pi_1} X_1 = \operatorname{Spec} R, \tag{10}$$

where $\pi_\mu : X_{\mu+1}\to X_\mu$ is the blowup of X_μ at the closed point $x_\mu\in X_\mu$ for every $\mu = 1,\dots,N$. We shall proceed by induction on N, the case $N=1$ being trivial. Consider the exceptional divisor E_N arising in the last blowup. Depending on whether x_N lies in an intersection of two exceptional divisor or not, E_N intersects one or two exceptional divisors. In other words, the vertex N is adjacent to one or two vertices.

Suppose first that the vertex N is adjacent to only one vertex β. We consider the ideal

$$\mathfrak{a}' := \mathfrak{p}_\beta^{\widehat{d}_N}\prod_{\nu\neq N}\mathfrak{p}_\nu^{\widehat{d}_\nu}.$$

It has the minimal principalization

$$X_N \xrightarrow{\pi_{N-1}} \cdots \xrightarrow{\pi_2} X_2 \xrightarrow{\pi_1} X_1 = \operatorname{Spec} R \tag{11}$$

The associated proximity matrix and its inverse are clearly restrictions of those of (10). So $k'_\nu = k_\nu$ for every $\nu\neq N$. Because N is adjacent only to β and $E_N^2=-1$, the proximity equation for the ideal $\mathfrak{p}_\nu$ gives $v_N(\mathfrak{p}_\nu) = v_\beta(\mathfrak{p}_\nu)$. By the reciprocity formula (7) we then obtain

$$v_\nu(\mathfrak{p}_N) = v_N(\mathfrak{p}_\nu) = v_\beta(\mathfrak{p}_\nu) = v_\nu(\mathfrak{p}_\beta).$$

Hence

$$d'_\nu = \widehat{d}_N v_\nu(\mathfrak{p}_\beta) + \sum_{\mu\neq N}\widehat{d}_\mu v_\nu(\mathfrak{p}_\mu) = \widehat{d}_N v_\nu(\mathfrak{p}_N) + \sum_{\mu\neq N}\widehat{d}_\mu v_\nu(\mathfrak{p}_\mu) = d_\nu$$

for all $\nu \neq N$. By the induction hypothesis, the claim therefore holds if $\gamma, \eta \neq N$. It remains to consider the numbers $\alpha_{\beta,N}$ and $\alpha_{N,\beta}$.

We first observe that N being proximate only to β, we have $q_{N,\nu} = q_{\beta,\nu}$ for all $\nu \neq N$ by (1). By (8) we thus get

$$k_N = \sum_{\nu} q_{N,\nu} = \sum_{\nu \neq N} q_{\beta,\nu} + 1 = k_\beta + 1.$$

Moreover, the base change formula (2) gives

$$\widehat{d}_N = \sum_{\nu} d_\nu(-E_\nu \cdot E_N) = d_N - d_\beta.$$

We then get

$$\alpha_{\beta,N} = k_\beta + 2 - \frac{k_\beta + 1}{d_\beta}(d_\beta + \widehat{d}_N) = 1 - \widehat{d}_N \frac{k_\beta + 1}{d_\beta} = 1 - \widehat{d}_N \frac{k_\beta + 1}{d'_\beta}.$$

Using (7), we have

$$v_\beta(\mathfrak{p}_\beta) = \sum_{\nu} q_{\beta,\nu}^2 \geq \sum_{\nu} q_{\beta,\nu} = k_\beta.$$

Thus

$$d'_\beta = \widehat{d}_N v_\beta(\mathfrak{p}_\beta) + \sum_{\nu \neq N} \widehat{d}_\nu v_\beta(\mathfrak{p}_\nu) \geq \widehat{d}_N k_\beta + \sum_{\nu \neq N} \widehat{d}_\nu v_\beta(\mathfrak{p}_\nu)$$

so that

$$0 < \widehat{d}_N \frac{k_\beta + 1}{d'_\beta} \leq \frac{k_\beta + 1}{k_\beta + \sum_{\nu \neq N} (\widehat{d}_\nu / \widehat{d}_N) v_\beta(\mathfrak{p}_\nu)} \leq \frac{k_\beta + 1}{k_\beta} \leq 2.$$

Therefore $-1 \leq \alpha_{\beta,N} < 1$. Moreover, by the above the equality $\alpha_{\beta,N} = -1$ can take place only if $k_\beta = 1$ and $\widehat{d}_\nu = 0$ for all $\nu \neq N$. This means that we must have $\beta = 1$ and $\mathfrak{a} = \mathfrak{p}_N^{\widehat{d}_N}$. In particular, N is the only vertex adjacent to β and $\widehat{d}_\beta = 0$. Thus a) holds for $\alpha_{\beta,N}$. In order to show that a) holds for $\alpha_{N,\beta}$, too, we note that

$$\alpha_{N,\beta} = -\alpha_{\beta,N} \frac{d_\beta}{d_N} = -\alpha_{\beta,N} \frac{d_\beta}{d_\beta + \widehat{d}_N}.$$

As necessarily $\widehat{d}_N > 0$, we see that $|\alpha_{N,\beta}| < 1$.

Let us then consider the case where the vertex N is adjacent to two vertices, say β and β'. Now let

$$\mathfrak{a}' := \mathfrak{p}_\beta^{\widehat{d}_N} \mathfrak{p}_{\beta'}^{\widehat{d}_N} \prod_{\nu \neq N} \mathfrak{p}_\nu^{\widehat{d}_\nu}.$$

The ideal $\mathfrak{a}'$ has a similar minimal principalization as in the preceeding case. Analogously one obtains $d'_\nu = d_\nu$, $k'_\nu = k_\nu$ for $\nu \neq N$, $d_N = d_\beta + d_{\beta'} + \widehat{d}_N$ and $k_N = k_\beta + k_{\beta'} + 1$. By induction we only need to prove the claim for $\alpha_{N,\beta}$, $\alpha_{N,\beta'}$ and $\alpha_{\beta,N}$, $\alpha_{\beta',N}$. We consider here only $\alpha_{N,\beta}$ and $\alpha_{\beta,N}$, the proof for $\alpha_{N,\beta'}$ and $\alpha_{\beta',N}$ being similar.

Set $m := v_\Gamma(\beta)$. Note that $\beta' \sim_{\Gamma'} \beta$, where Γ' denotes the dual graph of the minimal principalization of $\mathfrak{a}'$. Therefore we have $m = v_{\Gamma'}(\beta)$, too. By Lemma 3.2 we get

$$\sum_{\nu \sim_\Gamma \beta} \alpha_{\beta,\nu} = m - 2 + \widehat{d}_\beta \frac{k_\beta + 1}{d_\beta}$$

and

$$\sum_{\nu \sim_{\Gamma'} \beta} \alpha_{\beta,\nu} = m - 2 + (\widehat{d}_\beta + \widehat{d}_N) \frac{k_\beta + 1}{d_\beta}.$$

Together the above equations imply that

$$\alpha_{\beta,N} = \alpha_{\beta,\beta'} - \widehat{d}_N \frac{k_\beta + 1}{d_\beta}.$$

By the induction hypothesis, $\alpha_{\beta,\beta'} < 1$. Thereby also $\alpha_{\beta,N} < 1$. We still need to show that $\alpha_{\beta,N} \geq -1$. As $a_{\beta,\nu} < 1$ for every $N \neq \nu \sim \beta$, we see that

$$\sum_{N \neq \nu \sim_\Gamma \beta} \alpha_{\beta,\nu} \leq m - 1,$$

where the equality holds if and only if $m = 1$. Hence

$$\alpha_{\beta,N} = \sum_{\nu \sim_\Gamma \beta} \alpha_{\beta,\nu} - \sum_{N \neq \nu \sim_\Gamma \beta} \alpha_{\beta,\nu} \geq -1 + \widehat{d}_\beta \frac{k_\beta + 1}{d_\beta} \geq -1.$$

We conclude that $|\alpha_{\beta,N}| < 1$ unless $m = 1$ and $\widehat{d}_\beta = 0$, in which case $\alpha_{\beta,N} = -1$. Finally, we have

$$\alpha_{N,\beta} = -\alpha_{\beta,N} \frac{d_\beta}{d_N} = -\alpha_{\beta,N} \frac{d_\beta}{d_\beta + d_{\beta'} + \widehat{d}_N}.$$

Because $|\alpha_{\beta,N}| \leq 1$ and $\widehat{d}_N > 0$, we obtain $|\alpha_{N,\beta}| < 1$ as wanted.

b) Obviously, there is nothing to prove if γ has only one adjacent vertex. Thus we may assume that $m := v_\Gamma(\gamma) \geq 2$. Suppose that there are two vertices adjacent to γ, say η and η', with $\alpha_{\gamma,\eta}, \alpha_{\gamma,\eta'} \leq 0$. It then follows from Lemma 3.2 that

$$\sum_{\eta,\eta' \neq \nu \sim \gamma} \alpha_{\gamma,\nu} \geq m - 2.$$

On the other hand, $\alpha_{\gamma,\nu} < 1$ for every $\nu \sim \gamma$ by a). Subsequently, if $m > 2$, then

$$m - 2 > \sum_{\eta,\eta' \neq \nu \sim \gamma} \alpha_{\gamma,\nu}.$$

Hence necessarily $m = 2$, and further, by Lemma 3.2 we also have $\alpha_{\gamma,\eta} = 0 = \alpha_{\gamma,\eta'}$ and $\widehat{d}_\gamma = 0$. $\square$

Notation 3.4. *Let $\gamma \in |\Gamma|$ and let a_γ be a nonnegative integer. Suppose that*

$$\{\eta \mid \eta \sim \gamma\} = \{\eta_1, \dots, \eta_m\}.$$

Set

$$\Delta_j := d_{\eta_j} \lambda(a_\gamma, \gamma) - k_{\eta_j} - 1 \quad (j = 1, \dots, m)$$

so that

$$\lambda(\Delta_j, \eta_j) = \lambda(a_\gamma, \gamma).$$

Also write

$$\Delta_j = \lfloor \Delta_j \rfloor + \delta_j$$

where $0 \le \delta_j < 1$.

Lemma 3.5. *With the preceeding notation, we have*

$$\sum_{j=1}^{m} \Delta_j + \widehat{d}_\gamma \lambda(a_\gamma, \gamma) + m - 2 = w_\Gamma(\gamma) a_\gamma.$$

In particular,

$$\sum_{j=1}^{m} \lfloor \Delta_j \rfloor + m - 2 = w_\Gamma(\gamma) a_\gamma - \varphi,$$

where

$$\varphi := \sum_{j=1}^{m} \delta_j + \widehat{d}_\gamma \lambda(a_\gamma, \gamma)$$

is a nonnegative integer.

Proof. By the formulas (3) and (9)

$$\lambda(a_\gamma, \gamma) = \frac{w_\Gamma(\gamma)(a_\gamma + k_\gamma + 1)}{w_\Gamma(\gamma) d_\gamma} = \frac{w_\Gamma(\gamma) a_\gamma + \sum_{j=1}^{m} k_{\eta_j} + 2}{\sum_{j=1}^{m} d_{\eta_j} + \widehat{d}_\gamma}.$$

Therefore

$$\sum_{j=1}^{m} \Delta_j + \widehat{d}_\gamma \lambda(a_\gamma, \gamma) + m - 2 = w_\Gamma(\gamma) a_\gamma.$$

The last statement is now obvious. □

The following lemma, which can be considered as a generalization of Lemma 3.3, will play a crucial role in the sequel:

Lemma 3.6. *Given any vertex $\gamma \in |\Gamma|$ and any nonnegative integer a_γ, we may choose for every vertex $\eta \sim \gamma$ a nonnegative integer a_η so that*

$$w_\Gamma(\gamma) a_\gamma = \sum_{\eta \sim \gamma} a_\eta \quad \textit{and} \quad \lambda(a_\eta, \eta) \ge \lambda(a_\gamma, \gamma),$$

where the latter inequality holds for each $\eta \sim \gamma$ except at most one. More precisely, if

$$\{\eta \mid \eta \sim \gamma\} = \{\eta_1, \dots, \eta_m\},$$

where $m > 1$, then the following is true:

1) *If it is possible to find a nonnegative integer* a_{η_1} *with* $\lambda(a_{\eta_1}, \eta_1) = \lambda(a_\gamma, \gamma)$*, then one may choose the other integers* a_{η_j} *so that*

$$\lambda(a_{\eta_j}, \eta_j) \geq \lambda(a_\gamma, \gamma)$$

for all $1 < j \leq m$*. Moreover, this choice can be done in such a way that the strict inequality holds for all except at most one* $1 < j \leq m$*. In the case we already have a nonnegative integer* a_{η_2} *with* $\lambda(a_{\eta_2}, \eta_2) = \lambda(a_\gamma, \gamma)$*, we can assume that the inequality is strict for all* $2 < j \leq m$*.*

2) *If it is possible to find a nonnegative integer* a_{η_1} *with* $\lambda(a_{\eta_1}, \eta_1) < \lambda(a_\gamma, \gamma)$ *or, in the case* $\widehat{d}_\gamma > 0$*,* $\lambda(a_{\eta_1}, \eta_1) = \lambda(a_\gamma, \gamma)$*, then one can choose the other integers* a_η *in such a way that*

$$\lambda(a_{\eta_j}, \eta_j) > \lambda(a_\gamma, \gamma)$$

holds for every $1 < j \leq m$*.*

Proof. Obviously we may assume that $m > 1$.

In the case $a_\gamma = 0$, we set $a_\eta = 0$ for every $\eta \sim \gamma$. Since

$$\lambda(a_\eta, \eta) - \lambda(a_\gamma, \gamma) = \frac{\alpha_{\gamma,\eta}}{d_\eta},$$

the claim follows from Lemma 3.3.

Suppose thus that $a_\gamma > 0$. By Lemma 3.3 $\alpha_{\eta_j,\gamma} \geq -1$. Then

$$\alpha_{\gamma,\eta_j} = -\frac{d_{\eta_j}}{d_\gamma}\alpha_{\eta_j,\gamma} \leq \frac{d_{\eta_j}}{d_\gamma}.$$

As $a_\gamma > 0$, this implies that

$$\Delta_j = \frac{a_\gamma d_{\eta_j}}{d_\gamma} - \alpha_{\gamma,\eta_j} \geq 0.$$

By Lemma 3.5

$$\lfloor \Delta_1 \rfloor + \sum_{j=2}^{m} (\lfloor \Delta_j \rfloor + 1) = w_\Gamma(\gamma) a_\gamma + 1 - \varphi,$$

where

$$\varphi = \sum_{j=1}^{m} \delta_j + \widehat{d}_\gamma \lambda(a_\gamma, \gamma)$$

is a nonnegative integer.

We can assume that for some j there exists a nonnegative integer a_{η_j} with $\lambda(a_{\eta_j}, \eta_j) \leq \lambda(a_\gamma, \gamma)$. If this is not the case, then choose any nonnegative integers a_{η_j} satisfying

$$\sum_{j=1}^{m} a_{\eta_j} = w_\Gamma(\gamma) a_\gamma.$$

Then $\lambda(a_{\eta_j}, \eta_j) > \lambda(a_\gamma, \gamma)$ for all $j = 1, \ldots, m$ by the above assumption. But this means that we have proven the claim.

Suppose thus that, for example, $\lambda(a_{\eta_1}, \eta_1) \leq \lambda(a_\gamma, \gamma)$. We will first consider the case $\lambda(a_{\eta_1}, \eta_1) < \lambda(a_\gamma, \gamma)$. Because $\lambda(a_\gamma, \gamma) = \lambda(\Delta_{\eta_1}, \eta_1)$, this implies that $a_{\eta_1} < \Delta_1$. Then either $a_{\eta_1} \leq \lfloor \Delta_1 \rfloor - 1$ or $a_{\eta_1} = \lfloor \Delta_1 \rfloor$ and $\delta_1 > 0$. In the first case we write

$$\lfloor \Delta_1 \rfloor - 1 + \sum_{j=2}^{m} (\lfloor \Delta_j \rfloor + 1) = \sum_{j=1}^{m} \lfloor \Delta_j \rfloor + m - 2 \leq w_\Gamma(\gamma) a_\gamma,$$

whereas in the latter case we have $\varphi \geq 1$ so that

$$\lfloor \Delta_1 \rfloor + \sum_{j=2}^{m} (\lfloor \Delta_j \rfloor + 1) = \sum_{j=1}^{m} \lfloor \Delta_j \rfloor + m - 1 = w_\Gamma(\gamma) a_\gamma + 1 - \varphi \leq w_\Gamma(\gamma) a_\gamma.$$

It comes therefore out that it is possible to find numbers $a_{\eta_j} \geq \lfloor \Delta_j \rfloor + 1 > \Delta_j$ for $j = 2, \ldots, m$ such that

$$\sum_{j=1}^{m} a_{\eta_j} = w_\Gamma(\gamma) a_\gamma.$$

Because $\lambda(a_\gamma, \gamma) = \lambda(\Delta_j, \eta_j)$, $a_{\eta_j} > \Delta_j$ implies $\lambda(a_{\eta_j}, \eta_j) > \lambda(a_\gamma, \gamma)$.

Consider then the case $\lambda(a_{\eta_1}, \eta_1) = \lambda(a_\gamma, \gamma)$. Now $a_{\eta_1} = \Delta_1 = \lfloor \Delta_1 \rfloor$. We immediately observe that the above argument works if $\widehat{d_1} > 0$. This is the case also if some $\delta_j > 0$. We can therefore assume that $\delta_2 = 0$. Then $\Delta_2 = \lfloor \Delta_2 \rfloor$ is an integer. Take $a_{\eta_2} = \Delta_2$. We can now write

$$\lfloor \Delta_1 \rfloor + \lfloor \Delta_2 \rfloor + \sum_{j=3}^{m} (\lfloor \Delta_j \rfloor + 1) = \sum_{j=1}^{m} \lfloor \Delta_j \rfloor + m - 2 \leq w_\Gamma(\gamma) a_\gamma - \varphi \leq w_\Gamma(\gamma) a_\gamma.$$

Finally note that $\lambda(a_{\eta_2}, \eta_2) = \lambda(a_\gamma, \gamma)$ of course implies $a_{\eta_2} = \Delta_2$. $\square$

4. MAIN RESULTS

We first want to give a criterium for a positive rational number to be a jumping number. We begin with two lemmata. In the first one an antinef divisor is constructed for an ordered tree structure on the dual graph:

Lemma 4.1. *Let $S \subset |\Gamma|$ be a connected set of vertices. Suppose that there is a collection of nonnegative integers $\{a_\nu \in \mathbb{N} \mid d(\nu, S) \leq 1\}$ such that*

i) *$\lambda(a_\nu, \nu) > \xi = \lambda(a_\gamma, \gamma)$ for any $\gamma \in S$ and $\nu \sim S$;*

ii) *$w_\Gamma(\gamma) a_\gamma \geq \sum_{\nu \sim \gamma} a_\nu$ for every $\gamma \in S$.*

Then there exists an antinef divisor $F \in \Lambda$ such that

1) *$f_\nu = a_\nu$ for all $\nu \in |\Gamma|$ with $d(\nu, S) \leq 1$;*

2) *$\widehat{f_\nu} = 0$ for $\nu \notin S$ unless ν is an end;*

3) *For any $\nu \sim \mu$ such that $d(\nu, S) > d(\mu, S)$ we have*

$$\lambda(f_\nu, \nu) > \lambda(f_\mu, \mu).$$

Proof. If $N = 1$, then there is nothing to prove. Suppose thus that $N > 1$. We will define nonnegative integers f_ν inductively on $d(\nu, S)$. First set $f_\nu = a_\nu$ for all $\nu \in |\Gamma|$ with $d(\nu, S) \leq 1$. If $S = |\Gamma|$, we are done. Suppose then that f_ν has been defined for some $\nu \in |\Gamma|$ with $d(\nu, S) > 0$. Because S is connected and Γ contains no loops, there is a unique $\mu' \in |\Gamma|$ such that $\mu' \sim \nu$ and $d(\mu', S) = d(\nu, S) - 1$. By induction we know that $\lambda(f_\nu, \nu) > \lambda(f_{\mu'}, \mu')$. Therefore we can use Lemma 3.6 to find for each $\mu' \neq \mu \sim \nu$ a nonnegative integer f_μ such that $\lambda(f_\mu, \mu) > \lambda(f_\nu, \nu)$. In the case where ν is not an end we also get

$$\widehat{f_\nu} = w_\Gamma(\nu) f_\nu - \sum_{\mu \sim \nu} f_\mu = 0.$$

When all the numbers f_ν have so been defined, we can set $F = \sum_\nu f_\nu E_\nu$. It remains to show that F is antinef. This is equivalent to $\widehat{f_\nu} \geq 0$ for all ν. We have already seen that $\widehat{f_\nu} = 0$ if $\nu \notin S$ is not an end. Moreover, $\widehat{f_\nu} \geq 0$ when $\nu \in S$. In order to complete the proof we need to use the following Lemma 4.2. □

Lemma 4.2. *Assume that $N > 1$. Let τ be an end and μ the vertex adjacent to it. If $G \in \Lambda$ is such that $\lambda(g_\tau, \tau) > \lambda(g_\mu, \mu)$, then $\widehat{g}_\tau \geq 0$.*

Proof. Note first that $\widehat{g}_\tau = w_\Gamma(\tau) g_\tau - g_\mu$ by (3). The assumption now says that

$$\frac{g_\mu + k_\mu + 1}{d_\mu} < \frac{g_\tau + k_\tau + 1}{d_\tau}.$$

By the formulas (3) and (9)

$$\frac{w_\Gamma(\tau)\,(g_\tau + k_\tau + 1)}{w_\Gamma(\tau) d_\tau} = \frac{w_\Gamma(\tau) g_\tau + k_\mu + 2}{d_\mu + \widehat{d}_\tau} \leq \frac{(w_\Gamma(\tau) g_\tau + 1) + k_\mu + 1}{d_\mu}.$$

Thereby

$$\frac{g_\mu + k_\mu + 1}{d_\mu} < \frac{(w_\Gamma(\tau) g_\tau + 1) + k_\mu + 1}{d_\mu},$$

so that $g_\mu < w_\Gamma(\tau) g_\tau + 1$, i.e., $\widehat{g}_\tau \geq 0$. □

We are now able to prove our first main result:

Theorem 4.3. *Let $\mathfrak{a}$ be an ideal in a two-dimensional regular local ring R. A positive rational number ξ is a jumping number of $\mathfrak{a}$ if and only if there exists a connected set of vertices $S \subset |\Gamma|$ and a collection of nonnegative integers $\{a_\eta \in \mathbb{N} \mid d(\eta, S) \leq 1\}$ satisfying the following conditions:*

i) $\lambda(a_\eta, \eta) > \xi = \lambda(a_\gamma, \gamma)$ *for every $\gamma \in S$ and every $\eta \sim S$;*

ii) $w_\Gamma(\gamma) a_\gamma = \sum_{\nu \sim \gamma} a_\nu$ *for every vertex $\gamma \in S$ (when $N > 1$).*

The condition ii) can be replaced by the condition ii') $w_\Gamma(\gamma)a_\gamma \geq \sum_{\nu\sim\gamma} a_\nu$ *for every vertex* $\gamma \in S$.

When these conditions hold, there exists an antinef divisor F *with* $\xi = \xi_F$ *such that* $S = S_F$. *Finally, consider the conditions*

1) S *is a chain;*
2) *if* γ *is not an end of* S, *then* $\widehat{d}_\gamma = 0$*;*
3) *if* γ *is an end of* S, *then* γ *is a star or* $\widehat{d}_\gamma > 0$.

The set S *can then be chosen in such a way that conditions 1) and 2) hold while condition 3) is true for any* S.

Proof. The case $N = 1$ being trivial, we may assume that $N > 1$.

Suppose first that there exists a connected set $S \subset |\Gamma|$ and a collection of nonnegative integers $\{a_\nu \in \mathbb{N} | d(\nu, S) \leq 1\}$ satisfying conditions i) and ii'). Let $F \in \Lambda$ be an antinef divisor as in Lemma 4.1. For any chain $\gamma = \nu_0 \sim \cdots \sim \nu_r = \nu$ going away from S, where $\gamma \in S$, we then have

$$\lambda(f_\nu, \nu) > \cdots > \lambda(f_\gamma, \gamma) = \lambda(a_\nu, \nu) = \xi$$

so that $\xi = \min\{\lambda(f_\nu, \nu) | \nu \in |\Gamma|\}$. Then $\xi = \xi_F$ is a jumping number by Proposition 2.1. Note that for an end vertex γ of S, we have by Lemma 3.1 either $\widehat{d}_\gamma > 0$ or $v_\Gamma(\gamma) \geq 3$, which shows that condition 3) automatically holds for S.

Conversely, if ξ is a jumping number, then by Proposition 2.1 there exists an antinef divisor F such that $\xi = \xi_F$. We may now choose a connected component S of S_F and set $a_\nu := f_\nu$ for every ν with $d(\nu, S) \leq 1$. Clearly, the conditions i) and ii') are satisfied.

It remains to show that if there is a connected set $S' \subset |\Gamma|$ with a collection $\{a'_\nu \in \mathbb{N} | d(\nu, S') \leq 1\}$ that satisfies conditions i) and ii'), then we can find a new set $S \subset S'$ together with a collection $\{a_\nu \in \mathbb{N} | d(\nu, S) \leq 1\}$ satisfying conditions i) and ii). Moreover, also conditions 1) and 2) should hold for the set S.

If there is a vertex $\gamma \in S'$ which has only one adjacent vertex, say η, and $w_\Gamma(\gamma)a'_\gamma > a'_\eta$, then we choose $S := \{\gamma\}$ and take $a_\gamma := a'_\gamma$, $a_\eta := w_\Gamma(\gamma)a'_\gamma$.

When this is not the case, we look for chains consisting of vertices $\gamma \in S'$ such that $\widehat{d}_\gamma = 0$ if γ is not an end of the chain. We now take our set S to be any maximal chain of this type. Obviously conditions 1) and 2) then hold.

We will next define the integers a_ν for all $\nu \in |\Gamma|$ with $d(\nu, S) \leq 1$. To begin with, set $a_\nu := a'_\nu$ if $\nu \in S$. Then take any $\gamma \in S$ and look at the vertices $\eta \sim \gamma$. Note that if there is only one vertex η adjacent to γ, then we now necessarily have $w_\Gamma(\gamma)a'_\gamma = a'_\eta$. By the maximality of S, $\eta \in S'$ implies $\eta \in S$ so that $a_\eta := a'_\eta$ will do. Thus we may assume $v_\Gamma(\gamma) \geq 2$.

Suppose first that γ is an end of S such that $\lambda(a'_\eta, \eta) > \xi$ for all $\gamma \sim \eta \notin S$. By assumption ii') we have

$$w_\Gamma(\gamma)a_\gamma \geq \sum_{\gamma\sim\eta\in S} a_\eta + \sum_{\gamma\sim\eta\notin S} a'_\eta.$$

If the equality holds, we take $a_\eta := a'_\eta$ for every $\gamma \sim \eta \notin S$. When this is not the case, we may increase the a'_η:s, $\eta \notin S$, to find numbers a_η such that

$$w_\Gamma(\gamma)a_\gamma = \sum_{\eta\sim\gamma} a_\eta.$$

For any $\eta \sim \gamma$, $\eta \notin S$, we then have $\lambda(a_\eta, \eta) \geq \lambda(a'_\eta, \eta) > \xi$.

Otherwise, we can utilize Lemma 3.6 to define the numbers a_η. For this we note that if γ now is an end of S, then necessarily, by the maximality of S, we have $\widehat{d_\gamma} > 0$ and there must be a vertex $\eta_1 \in S$ adjacent to γ. Furthermore, when γ is not an end of S, there are two vertices $\eta_1, \eta_2 \in S$ adjacent to γ. ☐

Remark 4.4. *If* $\xi = \xi_F$ *where* F *is any antinef divisor, then it comes out from the proof of Theorem 4.3 that we can construct the set* S *in such a way that* $S \subset S_F$ *and that* $a_\gamma = f_\gamma$ *for all* $\gamma \in S$.

Recall that the first jumping number of $\mathfrak{a}$ is the log canonical threshold

$$\operatorname{lct}(\mathfrak{a}) = \min_\nu \frac{k_\nu + 1}{d_\nu}.$$

Corollary 4.5. *Consider the set of those vertices* $\gamma \in |\Gamma|$ *for which*

$$\operatorname{lct}(\mathfrak{a}) = \frac{k_\gamma + 1}{d_\gamma}.$$

This set then satisfies the conditions 1) – 3) of Theorem 4.3.

Proof. Recall first that $f_\gamma > 0$ for every $\gamma \in |\Gamma|$ if $0 \neq F \in \Lambda$ is an antinef divisor. The zero divisor 0 is then the unique antinef divisor $F \in \Lambda$ such that $\operatorname{lct}(\mathfrak{a}) = \xi_F$. Therefore the claim follows from Theorem 4.3. ☐

Remark 4.6. *This was observed by Veys in* [11, Theorem 3.3] *in the context of an embedded resolution of a curve.*

Example 4.7. *Conditions 1) and 2) of Theorem 4.3 do not hold for* S_F *for an arbitrary antinef divisor* $F \in \Lambda$. *To see this, consider the configuration of exceptional divisors described by the dual graph*



and the proximity matrix

$$P = \begin{bmatrix} 1 & 0 & 0 \\ {}^{-}1 & 1 & 0 \\ {}^{-}1 & 0 & 1 \end{bmatrix}.$$

The dual graph is then associated to a minimal principalization of the ideal $\mathfrak{a} = \mathfrak{p}_2\mathfrak{p}_3$. *Now*

$$D = 2E_1 + 3E_2 + 3E_3 = \widehat{E}_2 + \widehat{E}_3 \quad \text{and} \quad K = E_1 + 2E_2 + 2E_3.$$

Let us take

$$F := E_1 + E_2 + E_3 = \widehat{E}_1.$$

One now calculates that

$$\lambda(f_1, 1) = \frac{3}{2}, \lambda(f_2, 2) = \lambda(f_3, 3) = \frac{4}{3}.$$

Then $\xi_F = \frac{4}{3}$ *with* $S_F = \{2, 3\}$. *In particular, we see that* S_F *is disconnected. However, if*

$$F' := E_1 + E_2 + 2E_3 = \widehat{E}_3,$$

then also $\xi_{F'} = \frac{4}{3}$, *but* $S_{F'} = \{2\}$. *We now observe that* $S_{F'}$ *satisfies the conditions 1) – 3) of Theorem 4.3.*

Theorem 4.3 implies that a support of a jumping number always contains star vertices or vertices corresponding to Rees valuations. In Corollary 4.10 we are now going to show the converse: this kind of vertices always support some jumping number. First we need two lemmata:

Lemma 4.8. *Let* $\gamma \in |\Gamma|$. *If* $\nu_1 \prec \cdots \prec \nu_r = \eta$ *are points proximate to* γ, *then* $d_\eta > rd_\gamma$.

Proof. For every $i = 1, \ldots, r$, we have $\nu_i \succ \gamma$ and $\nu_i \succ \nu_{i-1}$, where $\nu_0 = \gamma$. Using the base change formula (2), we get

$$d_{\nu_i} = d_{\nu_{i-1}} + d_\gamma + d^*_{\nu_i} > d_{\nu_{i-1}} + d_\gamma$$

so that

$$d_\eta > d_{\nu_{r-1}} + d_\gamma > d_{\nu_{r-2}} + 2d_\gamma > \cdots > d_{\nu_1} + (r-1)d_\gamma > rd_\gamma.$$

□

Lemma 4.9. *Let* $\gamma \in |\Gamma|$.

a) *If* $\widehat{d}_\gamma > 0$, *then* $d_\gamma \geq k_\gamma \widehat{d}_\gamma$.
b) *If* $v_\Gamma(\gamma) + \widehat{d}_\gamma \geq 3$, *then* $d_\gamma - k_\gamma \geq 2$.

Proof. By (2) we have $d^*_\nu = \sum_\mu \widehat{d}_\mu q_{\mu,\nu} \geq \widehat{d}_\gamma q_{\gamma,\nu}$ for every $\nu \in |\Gamma|$. Again by (2), and (8), we then obtain

$$d_\gamma - k_\gamma \widehat{d}_\gamma = \sum_\nu (d^*_\nu - \widehat{d}_\gamma) q_{\gamma,\nu} \geq \widehat{d}_\gamma \sum_\nu (q_{\gamma,\nu} - 1) q_{\gamma,\nu} \geq 0$$

proving a).

In order to prove b), we first observe that $d_1^* = d_1 = v_1(\mathfrak{a}) \geq v_1(\mathfrak{p}_\gamma) = q_{\gamma,1} \geq 1$ and $d_1^* \geq d_\gamma^* \geq \widehat{d}_\gamma$. If now $q_{\gamma,1} \geq 2$ or if $\widehat{d}_\gamma \geq 3$, we get

$$d_\gamma - k_\gamma = \sum_\nu (d_\nu^* - 1) q_{\gamma,\nu} \geq (d_1^* - 1) q_{\gamma,1} \geq 2,$$

as wanted.

Consider then the case $q_{\gamma,1} = 1$ and $\widehat{d}_\gamma \leq 2$. The formula (1) implies that in this case γ is proximate to at most one adjacent vertex. The number of adjacent vertices proximate to γ must then be $v_\Gamma(\gamma) - t$, where $t \in \{0, 1\}$. By using (3), we get

$$d_\gamma^* = \widehat{d}_\gamma + \sum_{\nu \succ \gamma} d_\nu^* \geq \widehat{d}_\gamma + v_\Gamma(\gamma) - t \geq 3 - t.$$

Assuming $\gamma \neq 1$, i.e., $t = 1$, we now obtain

$$d_\gamma - k_\gamma \geq (d_1^* - 1) + (d_\gamma^* - 1) \geq 2.$$

In the case $\gamma = 1$ we have $t = 0$, and thus $d_\gamma^* \geq 3$. So $d_\gamma - k_\gamma \geq d_\gamma^* - 1 \geq 2$. □

We are now ready to prove the promised result:

Corollary 4.10. *Let* $\gamma \in |\Gamma|$.

a) *If* $\widehat{d}_\gamma$ *is positive, then*

$$\frac{n}{d_\gamma}$$

is a jumping number of $\mathfrak{a}$ *with a support* $\{\gamma\}$ *for any integer* n *such that* $n\widehat{d}_\gamma > d_\gamma$. *In particular, this implies that*

$$1 + \frac{n}{d_\gamma}$$

is a jumping number of $\mathfrak{a}$ *with a support* $\{\gamma\}$ *for every positive integer* n.

b) *If* $v_\Gamma(\gamma) + \widehat{d}_\gamma \geq 3$, *then*

$$1 - \frac{1}{d_\gamma}$$

is a jumping number of $\mathfrak{a}$ *with a support* $\{\gamma\}$. *This is the case especially when* γ *is a star.*

Proof. Suppose that $\{\eta \mid \eta \sim \gamma\} = \{\eta_1, \ldots, \eta_m\}$. Let a_γ be a nonnegative integer. By Lemma 3.5 we now have

$$\sum_{j=1}^m (\lfloor \Delta_j \rfloor + 1) = w_\Gamma(\gamma) a_\gamma + 2 - \varphi.$$

where

$$\varphi = \sum_{j=1}^m \delta_j + \widehat{d}_\gamma \lambda(a_\gamma, \gamma)$$

is a nonnegative integer. It follows that if $\varphi \geq 2$, then we can find nonnegative integers $a_{\eta_j} \geq \lfloor \Delta_j \rfloor + 1$ for $j = 1, \dots, m$ such that

$$\sum_{j=1}^{m} a_{\eta_j} = w_\Gamma(\gamma) a_\gamma$$

and that

$$\lambda(a_{\eta_j}, \eta_j) \geq \lambda(\lfloor \Delta_j \rfloor + 1, \eta_j) > \lambda(\Delta_j, \eta_j) = \lambda(a_\gamma, \gamma).$$

This means that we can use Theorem 4.3 to conclude that $\xi = \lambda(a_\gamma, \gamma)$ is a jumping number with a support $\{\gamma\}$. Let us now show that in both cases a) and b) we can choose a_γ in such a way that $\varphi \geq 2$.

Consider first the case a). For any integer n such that $n\widehat{d}_\gamma > d_\gamma$, set

$$a_\gamma = n - k_\gamma - 1.$$

By Lemma 4.9 a) a_γ is positive. Moreover

$$\lambda(a_\gamma, \gamma) = \frac{n}{d_\gamma}.$$

Then $\varphi \geq \left\lceil \widehat{d}_\gamma \lambda(a_\gamma, \gamma) \right\rceil \geq 2$ as wanted.

Consider then the case b). Set $a_\gamma = d_\gamma - k_\gamma - 2$. Now Lemma 4.9 b) implies that a_γ is nonnegative. Furthermore,

$$\lambda(a_\gamma, \gamma) = 1 - \frac{1}{d_\gamma}.$$

Then

$$\Delta_j = d_{\eta_j} \lambda(a_\gamma, \gamma) - k_{\eta_j} - 1 = -\frac{d_{\eta_j}}{d_\gamma} + d_{\eta_j} - k_{\eta_j} - 1.$$

for all $j = 1, \dots, m$. Thus

$$\delta_j = \Delta_j - \lfloor \Delta_j \rfloor = \left\lceil \frac{d_{\eta_j}}{d_\gamma} \right\rceil - \frac{d_{\eta_j}}{d_\gamma}.$$

By the formula (3)

$$w_\Gamma(\gamma) d_\gamma = \sum_{j=1}^{m} d_{\eta_j} + \widehat{d}_\gamma.$$

Hence

$$\sum_{j=1}^{m} \delta_j = \sum_{j=1}^{m} \left\lceil \frac{d_{\eta_j}}{d_\gamma} \right\rceil - \sum_{j=1}^{m} \frac{d_{\eta_j}}{d_\gamma} = \sum_{j=1}^{m} \left\lceil \frac{d_{\eta_j}}{d_\gamma} \right\rceil - w_\Gamma(\gamma) + \frac{\widehat{d}_\gamma}{d_\gamma}$$

so that

$$\varphi = \sum_{j=1}^{m} \delta_j + \widehat{d}_\gamma \lambda(a_\gamma, \gamma) = \sum_{j=1}^{m} \left\lceil \frac{d_{\eta_j}}{d_\gamma} \right\rceil - w_\Gamma(\gamma) + \widehat{d}_\gamma.$$

Since $\gamma \sim \eta_j$, we have either $\gamma \prec \eta_j$ or $\eta_j \prec \gamma$. In the first case we have a maximal chain $\nu_1 \prec \cdots \prec \nu_{r_j} = \eta_j$ of points proximate to γ. By Lemma 4.8 this means that $d_{\eta_j} \geq r_j d_\gamma + 1$. This certainly holds also when $\eta_j \prec \gamma$ if we set $r_j = 0$ in this case. Therefore we have, for every $j = 1, \dots, m$,

$$\left\lceil \frac{d_{\eta_j}}{d_\gamma} \right\rceil \geq 1 + r_j \quad \text{and} \quad \sum_{j=1}^{m} r_j = \#\{\nu \in |\Gamma| \mid \nu \succ \gamma\} = w_\Gamma(\gamma) - 1.$$

It follows that

$$\sum_{j=1}^{m} \left\lceil \frac{d_{\eta_j}}{d_\gamma} \right\rceil \geq m - 1 + w_\Gamma(\gamma),$$

and further, $\varphi \geq m - 1 + \widehat{d_\gamma} \geq 2$, which proves the claim. □

Remark 4.11. *In the case of a curve on a smooth surface, it was observed by Smith and Thompson in* [9, Theorem 3.1] *that b) of Corollary 4.10 holds for any star vertex.*

We will next show how new jumping nunbers can be obtained from a given one.

Proposition 4.12. *Suppose ξ is a jumping number of $\mathfrak{a}$ with a support $S \subset |\Gamma|$. Write*

$$d = \gcd\{d_\eta | d(\eta, S) \leq 1\}.$$

Then, for any $n \in \mathbb{N}$,

$$\xi + \frac{n}{d}$$

is also a jumping number of $\mathfrak{a}$ with a support S.

Proof. Suppose first that S is connected. Let $\{a_\eta | d(\eta, S) \leq 1\}$ be a collection of nonnegative integers satisfying the conditions of Theorem 4.3. For every vertex $\eta \in |\Gamma|$ with $d(\eta, S) \leq 1$, write

$$b_\eta := \frac{d_\eta}{d}.$$

Clearly,

$$\lambda(a_\eta + nb_\eta, \eta) = \lambda(a_\eta, \eta) + \frac{n}{d}$$

for any $\eta \in |\Gamma|$ and $n \in \mathbb{N}$. Hence

$$\lambda(a_\eta + nb_\eta, \eta) > \xi + \frac{n}{d} = \lambda(a_\gamma + nb_\gamma, \gamma)$$

for every $\gamma \in S$ and $\eta \sim S$. By the proximity inequality for $\mathfrak{a}$

$$w_\gamma b_\gamma d = w_\gamma d_\gamma \geq \sum_{\nu \sim \gamma} d_\nu = d \sum_{\nu \sim \gamma} b_\nu.$$

We thus see that

$$w_\gamma(a_\gamma + nb_\gamma) \geq \sum_{\nu \sim \gamma} (a_\nu + nb_\nu).$$

for every $\gamma \in S$. The claim then results from Theorem 4.3.

Consider then the general case. Let $S_1, \ldots, S_r$ be the connected components of S. Write

$$d_i := \gcd\{d_\eta | d(\eta, S_i) \leq 1\} \quad (i = 1, \ldots, r).$$

Then clearly $d = \gcd\{d_1, \ldots, d_r\}$. By the preceeding case we now know that

$$\xi + \frac{n}{d}$$

is a jumping number with a support S_i for every $i = 1, \ldots, r$. Then the claim follows from the following Lemma 4.13:

$\square$

Lemma 4.13. *If $S_1, S_2 \subset |\Gamma|$ are supports of a jumping number ξ, then so is $S_1 \cup S_2$.*

Proof. Let $F_1, F_2 \in \Lambda$ be antinef divisors such that $S_1 = S_{F_1}$ and $S_2 = S_{F_2}$. Let us define a divisor F by setting

$$f_\nu := \min\{f_{1,\nu}, f_{2,\nu}\}.$$

Since F_1 and F_2 are both antinef, we see that

$$\sum_{\eta \sim \gamma} f_\eta \leq \sum_{\eta \sim \gamma} f_{i,\eta} \leq w_\Gamma(\gamma) f_{i,\gamma}$$

for any $\gamma \in |\Gamma|$ and $i = 1, 2$, and further,

$$\sum_{\eta \sim \gamma} f_\eta \leq w_\Gamma(\gamma) f_\gamma.$$

Hence F is antinef, too. Clearly, $\lambda(f_\nu, \nu) \geq \xi$, where the equality holds if and only if $\nu \in S_1 \cup S_2$. Therefore $S_1 \cup S_2 = S_F$, which proves the claim. $\square$

Tucker introduced in [10] the notions of contibution and critical contribution of a jumping number by a divisor. The contribution of a prime divisor had earlier been defined by Smith and Thompson in [9]. Consider a reduced subdivisor

$$G = E_{\gamma_1} + \ldots + E_{\gamma_r} \leq D.$$

One first says that a positive rational number ξ is a *candidate jumping number* for G if ξd_γ is an integer for all $\gamma \in \{\gamma_1, \ldots, \gamma_r\}$. The divisor G is then said to *contribute* the jumping number ξ if ξ is a candidate jumping number for G and

$$\mathcal{J}(\mathfrak{a}^\xi) \subsetneq \Gamma(X, \mathcal{O}_X(G + K - \lfloor \xi D \rfloor)).$$

Finally, a contribution is said to be *critical* if, in addition, no proper subdivisor of G contributes ξ, i. e.,

$$\mathcal{J}(\mathfrak{a}^\xi) = \Gamma(X, \mathcal{O}_X(G' + K - \lfloor \xi D \rfloor))$$

for all divisors $0 \leq G' < G$.

Proposition 4.14. *If a divisor G critically contributes ξ, then $\xi = \xi_F$ for an antinef divisor F with $S_F = |G|$.*

Proof. Set $F := (\lfloor \xi D \rfloor - K - G \rfloor)^{\sim}$. Because ξ is a candidate jumping number for G, we have

$$G + K - \lfloor \xi D \rfloor \leq K - \lfloor (\xi - \epsilon) D \rfloor$$

for small enough $\epsilon > 0$. Let $\mathfrak{b} = \Gamma(X, \mathcal{O}(-F))$ be the complete ideal associated to F. By the above we get $\mathfrak{b} \subset \mathcal{J}(\mathfrak{a}^{\xi-\epsilon})$. Clearly $\mathcal{J}(\mathfrak{a}^{\xi}) \subset \mathfrak{b}$, but as G is contributing, we must have $\mathcal{J}(\mathfrak{a}^{\xi}) \neq \mathfrak{b}$. It then follows from Proposition 2.1 that $\xi = \xi_F$. Let us now show that $\gamma \in |G|$ is equivalent to $f_\gamma = \xi d_\gamma - k_\gamma - 1$. Since $\xi = \xi_F$, we have

$$f_\gamma \geq \xi d_\gamma - k_\gamma - 1$$

for all $\gamma \in |\Gamma|$. Now $\lfloor \xi D \rfloor - K - G \leq F$. For $\gamma \notin |G|$, this means that

$$f_\gamma \geq \lfloor \xi d_\gamma \rfloor - k_\gamma > \xi d_\gamma - k_\gamma - 1.$$

Suppose that we would have $f_\gamma \geq \xi d_\gamma - k_\gamma$ for some $\gamma \in |G|$. Write $G' := G - E_\gamma$. Then G' is a proper subdivisor of G and

$$\lfloor \xi D \rfloor - K - G \leq \lfloor \xi D \rfloor - K - G' \leq F = (\lfloor \xi D \rfloor - K - G)^{\sim}$$

implying that $(\lfloor \xi D \rfloor - K - G')^{\sim} = F$. But then

$$\mathcal{J}(\mathfrak{a}^{\xi}) \subsetneq \Gamma(X, \mathcal{O}(K + G' - \lfloor \xi D \rfloor))$$

which is impossible, because G critically contributes ξ. □

Lemma 4.15. *Suppose that F is an antinef divisor such that $\xi = \xi_F$. Then*

$$G := \sum_{\gamma \in S_F} E_\gamma$$

is a contributing divisor for ξ.

Proof. Because $\lambda(f_\gamma, \gamma) = \xi$ for all $\gamma \in S_F$, ξ is a candidate jumping number for G. Now

$$f_\nu \geq \xi d_\nu - k_\nu - 1$$

where the equality holds if and only if $\nu \in S_F$. In other words,

$$f_\nu \geq \lfloor \xi d_\nu \rfloor - k_\nu$$

if $\nu \notin S_F$ whereas

$$f_\nu = \lfloor \xi d_\nu \rfloor - k_\nu - 1$$

if $\nu \in S_F$. But this says that

$$-F \leq G + K - \lfloor \xi D \rfloor$$

implying that

$$\Gamma(X, \mathcal{O}(-F)) \subset \Gamma(X, \mathcal{O}_X(G + K - \lfloor \xi D \rfloor)).$$

Because $\xi = \xi_F$, it then follows from Proposition 2.1 that we cannot have

$$\Gamma(X, \mathcal{O}_X(G + K - \lfloor \xi D \rfloor)) = \mathcal{J}(\mathfrak{a}^{\xi}).$$

□

The following theorem now connects our approach to that of Smith, Thompson and Tucker:

Theorem 4.16. *Let $\mathfrak{a}$ be an ideal in a two-dimensional regular local ring R. Let G be a divisor critically contributing a jumping number ξ of $\mathfrak{a}$. Then there exists an antinef divisor F such that $\widehat{f_\gamma} = 0$ unless γ is an end not in $|G|$, $\lambda(a_\gamma, \gamma)$ reaches its minimum ξ exactly when $\gamma \in |G|$ and grows strictly on every path away from $|G|$. Moreover,*

1) *$|G|$ is a chain;*
2) *if $\gamma \in |G|$ has more than one adjacent vertex in $|G|$ then $\widehat{d_\gamma} = 0$;*
3) *if γ is an end of $|G|$ then γ is a star or $\widehat{d_\gamma} > 0$.*

Proof. According to Proposition 4.14 there is an antinef divisor F' such that $\lambda(f'_\gamma, \gamma) = \xi$ exactly when $\gamma \in |G|$. Let F be an antinef divisor as in Theorem 4.3 such that

$$S_F := \{\gamma \in \Gamma \mid \lambda(f_\gamma, \gamma) = \xi\} \subset \{\gamma \in \Gamma \mid \lambda(f'_\gamma, \gamma) = \xi\} = |G|$$

(see Remark 4.4). By Lemma 4.15 the divisor $\sum_{\nu \in S_F} E_\nu$ is contributing. As G is critically contributing, this implies that $S_F = |G|$, and we are done. □

Remark 4.17. *It was discovered by Tucker in* [10, Theorem 5.1] *that if G is a divisor critically contributing a jumping number, then $|G|$ satisfies conditions 1) – 3).*

Unfortunately, the existence of an antinef divisor F as in Theorem 4.16 does not guarantee that the contribution is critical. This will come out from the following example:

Example 4.18. *Consider the configuration of exceptional divisors described by the dual graph*



and the proximity matrix

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ {}^{-}1 & 1 & 0 & 0 & 0 \\ {}^{-}1 & 0 & 1 & 0 & 0 \\ {}^{-}1 & 0 & 0 & 1 & 0 \\ {}^{-}1 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

They are associated to the minimal principalization of the ideal $\mathfrak{a} = \mathfrak{p}_2^4\mathfrak{p}_3^4\mathfrak{p}_4^5\mathfrak{p}_5^7$. *Now*

$$D = 20E_1 + 24E_2 + 24E_3 + 25E_4 + 27E_5$$

and

$$K = E_1 + 2E_2 + 2E_3 + 2E_4 + 2E_5.$$

Let us take

$$F := 3E_1 + 3E_2 + 3E_3 + 4E_4 + 5E_5 = \widehat{E}_4 + 2\widehat{E}_5.$$

One then calculates that

$$\lambda(f_1, 1) = \lambda(f_2, 2) = \lambda(f_3, 3) = \frac{1}{4}, \lambda(f_4, 4) = \frac{7}{25}, \lambda(f_5, 5) = \frac{8}{27}.$$

Thus $\xi_F = \frac{1}{4}$ *and* $S_F = \{1, 2, 3\}$. *Set* $G = E_1 + E_2 + E_3$. *We now observe that the divisor* F *satisfies the conditions of Theorem 4.16. However,* G *is not critically contributing. Indeed, set*

$$F' := 3E_1 + 4E_2 + 3E_3 + 4E_4 + 4E_5 = \widehat{E}_2 + \widehat{E}_4 + \widehat{E}_5$$

and

$$F'' := 3E_1 + 3E_2 + 4E_3 + 4E_4 + 4E_5 = \widehat{E}_3 + \widehat{E}_4 + \widehat{E}_5.$$

Then

$$\lambda(f'_1, 1) = \frac{5}{20}, \lambda(f'_2, 2) = \frac{7}{24}, \lambda(f'_3, 3) = \frac{6}{24}, \lambda(f'_4, 4) = \frac{7}{25}, \lambda(f'_5, 5) = \frac{7}{27}$$

and

$$\lambda(f''_1, 1) = \frac{5}{20}, \lambda(f''_2, 2) = \frac{6}{24}, \lambda(f''_3, 3) = \frac{7}{24}, \lambda(f''_4, 4) = \frac{7}{25}, \lambda(f''_5, 5) = \frac{7}{27}.$$

Therefore $\xi_{F'} = \xi = \xi_{F''}$ *but*

$$S_{F'} = \{1, 3\} \subsetneq S_F \supsetneq \{1, 2\} = S_{F''}.$$

By Lemma 4.15 both $S_{F'}$ *and* $S_{F''}$ *contain subsets corresponding to divisors contributing* ξ*, but then the divisor* G *cannot be critically contributing. Moreover, it is easy to see that both* $E_1 + E_2$ *and* $E_1 + E_3$ *critically contribute* ξ*. Note that their intersection* E_1 *does not contribute* ξ*. Observe also that in this example the set* S_F *includes a star vertex which is a non-end of* S_F*.*

We next give a combinatorical criterium for the critical contribution:

Theorem 4.19. *Let $\mathfrak{a}$ be an ideal in a two-dimensional regular local ring R. Let G be a reduced subdivisor of D such that $|G|$ contains at least two vertices. Then G critically contributes a jumping number ξ if and only if $|G|$ is connected and there exists a collection of nonnegative integers $\{a_\nu \in \mathbb{N} \mid d(\nu, |G|) \leq 1\}$ such that*

i) $\lambda(a_\nu, \nu) > \xi = \lambda(a_\gamma, \gamma)$ *for any* $\gamma \in |G|$ *and* $\nu \sim |G|$*;*

ii) $w_\Gamma(\gamma) a_\gamma = \sum_{\nu \sim \gamma} a_\nu$ *for every* $\gamma \in |G|$*;*

iii) $\lambda(a_\nu - 1, \nu) \leq \xi$ *for any vertex* $\nu \sim |G|$*.*

Moreover, it is enough that iii) holds for vertices $\nu \sim \gamma$, where $\gamma \in |G|$ is a star or $\widehat{d}_\gamma > 0$.

Finally, if G critically contributes ξ and F is any antinef divisor such that ξ has support $|G|$ with respect to F, then we may take $a_\nu = f_\nu$ for all ν with $d(\nu, |G|) \leq 1$.

Proof. Suppose first that $|G|$ is connected and that there is a collection of nonnegative integers $\{a_\nu \in \mathbb{N} \mid d(\nu, |G|) \leq 1\}$ satisfying conditions i) – iii). Let $F \in \Lambda$ be an antinef divisor as in Lemma 4.1. This means that

$$S_F = \{\gamma \in |\Gamma| \mid \lambda(f_\gamma, \gamma) = \xi\} = |G|.$$

It then follows from Lemma 4.15 that the divisor G is contributing. For some subset $S \subset |G|$ the divisor $\sum_{\nu \in S} E_\nu$ is necessarily critically contributing. If $S \neq |G|$, then there must be a vertex $\eta \in |G| \smallsetminus S$ adjacent to a vertex $\gamma \in S$. By Theorem 4.16 we can find an antinef divisor F' such that $\xi = \xi_{F'}$ and $S = S_{F'}$. Then $\lambda(f'_\eta, \eta) > \xi = \lambda(a_\eta, \eta)$ so that $f'_\eta > a_\eta$. Because $\widehat{f'_\gamma} = 0$, we have

$$\sum_{\nu \sim \gamma} f'_\nu = w_\Gamma(\gamma) f'_\gamma = w_\Gamma(\gamma) a_\gamma = \sum_{\nu \sim \gamma} a_\nu.$$

Thus there must be at least one vertex $\mu \sim \gamma$ for which $f'_\mu < a_\mu$. So $\lambda(f'_\mu, \mu) < \lambda(a_\mu, \mu)$. This implies $\mu \notin |G|$ as otherwise $\lambda(f'_\mu, \mu) < \xi$ which is impossible. If γ is adjacent to some vertex in S, then γ must be star. If γ is not a star, then by Theorem 4.16 $S = \{\gamma\}$ and $\widehat{d}_\gamma > 0$. Condition iii) now implies that

$$\lambda(f'_\mu, \mu) \leq \lambda(a_\mu - 1, \mu) \leq \xi.$$

However, this is possible only if $\lambda(f'_\mu, \mu) = \xi$. But this means that $\mu \in S$, which is a contradiction. Therefore we must have $S = |G|$ as wanted.

Suppose then that G critically contributes ξ. Let $\gamma \in |G|$. By Theorem 4.14 there then exists an antinef divisor F such that $\lambda(f_\nu, \nu) > \xi = \lambda(f_\gamma, \gamma)$ for any $\nu \notin |G|$. Set $a_\nu = f_\nu$ for all $\nu \in |\Gamma|$ with $d(\nu, |G|) \leq 1$. Suppose that there would exist a vertex $\eta \sim \gamma$ such that $\lambda(a_\eta - 1, \eta) > \lambda(a_\gamma, \gamma)$. We may choose a vertex $\gamma' \in |G|$ such that $\gamma' \sim \gamma$. Let S be a connected component of $|G| \setminus \{\gamma'\}$ containing γ. Choose a collection of nonnegative integers $\{a'_\nu \in \mathbb{N} \mid d(\nu, S) \leq 1\}$ so that $a'_\eta = a_\eta - 1$ and

$a'_{\gamma'} = a_{\gamma'} + 1$ while $a'_\nu = a_\nu$ otherwise. Clearly $\lambda(a'_\nu, \nu) > \xi$ for any $\nu \sim S$. Also

$$w_\Gamma(\gamma) a'_\mu = \sum_{\nu \sim \mu} a'_\nu$$

for every $\mu \in S$. Therefore we can use Lemma 4.1 to find an antinef divisor F' such that $S = S_{F'}$. By Lemma 4.15 the divisor $\sum_{\nu \in S} E_\nu$ is then contributing. But this is a contradiction, because G is critically contributing. □

Example 4.20. *Consider the situation of Example 4.18. We now have*

$$\lambda(f_5 - 1, 5) = \frac{7}{27} > \frac{1}{4}.$$

It therefore follows from Theorem 4.19 that $E_1 + E_2 + E_3$ *does not critically contribute* ξ*.*

References

[1] A. Campillo, G. Gonzales-Sprinberg and M. Lejeune-Jalabert, *Clusters of infinitely near points*, Math. Ann. **306** (1996), 169–194.

[2] L. Ein, R. Lazarsfeld, K. Smith and D. Varolin, *Jumping coefficients of multiplier ideals*, Duke Math. J. **123** (2004), no. 3, 469–506.

[3] C. Favre, M. Jonsson, *The valuative tree*, Lecture Notes in Mathematics **1853**, Springer-Verlag, Berlin, 2004.

[4] J. Lipman, *Proximity inequalities for complete ideals in two-dimensional regular local rings*, Contemp. Math. **159** (1994), 293–306.

[5] J. Lipman and K. Watanabe, *Integrally closed ideals in two-dimensional regular local rings are multiplier ideals*, Math. Res. Lett. **10** (2003), no. 4, 423–434.

[6] F. Loeser, *Fonctions d'Igusa p-adiques et polynômes de Bernstein*, Amer. J. Math. **110** (1988), 1–21.

[7] T. Järvilehto, *Jumping numbers of a simple complete ideal in a two-dimensional regular local ring*, to appear in Mem. Am. Math. Soc.

[8] L. Van Proeyen and W. Veys, *Poles of the topological zeta function associated to an ideal in dimension two*, Math. Z. **260** (2008), no. 3, 615–627.

[9] K. Smith and H. Thompson, *Irrelevant exceptional divisors for curves on a smooth surface*, Algebra, geometry and their interactions, Contemp. Math. **448** (2007), 245–254.

[10] K. Tucker, *Jumping numbers on algebraic surfaces with rational singularities*, Trans. Amer. Math. Soc. **362** (2010), no. 6, 3223-3241.

[11] W. Veys, *Determination of the poles of the topological zeta function for curves*, Manuscripta Math. **87** (1995), no. 4, 435–448.

Mathematics and Statistics, School of Information Sciences, University of Tampere, FIN-33014 Tampereen yliopisto, Finland
E-mail address: `eero.hyry@uta.fi`

Pääskykuja 5, FIN-04620 Mäntsälä, Finland
E-mail address: `tarmo.jarvilehto@helsinki.fi`